

Not for Sale, Sale Into, Distribution and/or Use in Nassau, Suffolk, Kings, Queens Counties of New York State



GROUP 28 INSECTICIDE

Acelepryn® G

syngenta®

Insecticide

For systemic control of white grubs and other listed pests infesting landscape and recreational turfgrass (including golf courses) as well as landscape ornamentals, interior plantscapes and sod farms.

Active Ingredient:

Chlorantraniliprole¹

3-Bromo-N-[4-chloro-2-methyl-6-[(methylamino)carbonyl]phenyl]

-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxamide 0.2%

Other Ingredients 99.8%

Total: 100.0%

¹Chlorantraniliprole belongs to the anthranilic diamide chemical class

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 100-1500 EPA Est. No. 67572-GA-1

Product of USA

PRODUCT ID.

53199

25 pounds
Non-refillable Container

TM

Acelepryn® G

FIRST AID

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),
Call
1-800-888-8372

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

When used as directed, this product does not present a hazard to humans or domestic animals.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

After the product has been loaded into properly calibrated application equipment, shirt, pants, socks and shoes are sufficient Personal Protective Equipment. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates, oysters and shrimp. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to use sites.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and understand the entire label before using this product.

Acelepryn® G must be used only in accordance with specifications on this label or in separate Syngenta supplemental labeling that may be made temporarily available through local distributors, as a result of new EPA approvals. Syngenta will not be responsible for losses or damages resulting from the use of this product in any manner not specifically stated on this label or other labels or bulletins published by Syngenta. User assumes all risks associated with such non-specified use.

Acelepryn® G

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Professional applications to golf courses, residential, industrial and commercial lawns and sports fields are not within the scope of the Worker Protection Standard.

Do not allow other workers to enter the treated area while application is in progress. Do not apply this product in commercial nurseries and greenhouses.

PRODUCT INFORMATION

Acelepryn G may be applied for the control of white grubs and other listed pests infesting landscape and recreational turfgrass (including golf courses) as well as landscape ornamentals and interior plantscapes.

Acelepryn G may be used on plants or turfgrasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns, commercial lawns, industrial facilities, residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes, institutional buildings, airports, cemeteries; interior plantscapes, ornamental gardens, wildlife plantings, parks, playgrounds, schools, day-care facilities, golf courses (tee box areas, roughs, fairways, greens, collars etc.), athletic fields, sod farms and other landscaped areas.

Do not apply Acelepryn G to soil saturated with water because adequate distribution of the active ingredient vertically in the soil profile cannot be achieved under this condition.

Acelepryn G is in GROUP 28 of the EPA's Insecticide and Acaricide Groups Based on Target Site of Action (EPA PR Notice 2001-5) and may be used in rotational resistance management programs.

Consult your local Syngenta representative, Cooperative Extension Service specialist or pest control advisor for regionally specific information regarding application timing.

APPLICATION EQUIPMENT

Apply Acelepryn G with drop-type, rotary-type or hand-held application equipment. The application equipment must be properly calibrated according to the manufacturer's directions to ensure uniform coverage over the treated area. During application, check equipment periodically to confirm that it is working properly and that the granules are being uniformly distributed.

APPLICATION TIMING

White Grubs: Apply Acelepryn G from early April to early September for preventative and early curative control of all the major white grub species infesting turfgrass. The need for an application may be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. The higher rate listed for white grub control in Table 1 may be required in late August or early September when less sensitive mid-instar grubs are present at the time of application. Irrigate turf immediately after application or allow rainfall to move the product into the soil.

Annual Bluegrass Weevil: Apply Acelepryn G when overwintered adult annual bluegrass weevils are observed in late April or early May to prevent damage from first generation larvae in late-May and June. An application of Acelepryn G at this time will also provide white grub control.

Billbugs: Apply Acelepryn G when overwintered adult billbugs are first observed. This will usually occur in late April or early May in regions with cool-season turfgrasses. An application of Acelepryn G at this time will also provide white grub control. The higher rates listed in Table 1 may be required for applications made after early May or when billbug species other than bluegrass billbug are present.

Acelepryn® G

European Crane Fly: Apply Acelepryn G between September and November to control European crane fly larvae in turfgrass. The higher rate listed in Table 1 may be required to achieve control when applications are made in November.

Turf Caterpillars: Acelepryn G will provide curative caterpillar control in turfgrass. If the area being treated is maintained at a mowing height of greater than one inch, then the higher rate listed in Table 1 may be required during periods of high pest pressure.

Chinch Bugs: For suppression of chinch bugs, apply Acelepryn G before eggs hatch.

APPLICATION RATES FOR LAWNS, GOLF COURSES AND OTHER RECREATIONAL TURFGRASS AREAS

Table 1 provides the application rates that will control the listed turfgrass pests. For maximum residual effectiveness or for optimal control of large pest infestations, Acelepryn G may be applied at up to 125 pounds per acre (0.25 lb AI/A) to control any of the pests listed in Table 1. Restriction: Do not apply more than 250 pounds (equivalent to 0.5 lb of active ingredient) of product per acre per year in broadcast applications to turfgrass.

Table 1: Turfgrass Application Rates

Target Pest	Product per Acre	Product per 1,000 sq ft	Lb AI/A
White Grubs (including <i>Aphoditus</i> spp., Asiatic garden beetle, black turfgrass ataeenius, European chafer, green June beetle, Japanese beetle, May/June beetles (<i>Phyllophaga</i> spp.), northern masked chafer, oriental beetle and southern masked chafer) Billbugs European Crane Fly Turf caterpillars (including armyworms, cutworms and sod worms)	50 to 100 lb	1.15 to 2.3 lb	0.1 to 0.2
Annual Bluegrass weevil	75 to 100 lb	1.72 to 2.3 lb	0.15 to 0.2
Chinch bugs (suppression only)	50 to 100 lb	1.15 to 2.3 lb	0.1 to 0.2

APPLICATION RATES FOR ORNAMENTAL PLANTS

Table 2 provides the application rates that will control the pests of landscape ornamentals and interior plantscapes. Before application, remove black plastic or other weed barrier materials that are impermeable to water. For maximum residual effectiveness or for optimal control of large pest infestations, Acelepryn G may be applied at up to 125 pounds per acre (0.25 lb AI/A) to control any of the pests listed in Table 2.

Table 2: Ornamental Tree and Shrub (Including Evergreens), Flower, Foliage Plant, and Groundcover Application Rates

Target Pest	Product per Acre	Product per 1,000 sq ft	Lb AI/A
White Grubs (including <i>Aphoditus</i> spp., Asiatic garden beetle, black turfgrass Ataeenius, European chafer, green June beetle, Japanese beetle, May/June beetles (<i>Phyllophaga</i> spp.), northern masked chafer, oriental beetle and southern masked chafer)	50 to 100 lb	1.15 to 2.3 lb	0.1 to 0.2

Acelepryn® G

Table 3: Spreader Settings for These Application Rates

Manufacturer	Model	Effective Swath Width (Feet)	50 lb per Acre	75 lb per Acre	100 lb per Acre	125 lb per Acre
"Vicon"	03	20	11	12	14	16
"Lely"	W	34	3 1/2	4	4 1/4	4 3/4
"Scotts"	R8A	11 (Pattern 5)	H	H 1/2	I 1/2	J 1/4
Andersons	"Accupro 2000"	11 (Pattern 6)	I 1/2	J	J 3/4	K 1/2
"LESCO"	020093	5 (Pattern 1/4 Closed)	C	C 1/4	C 3/4	D 1/4
"Prizelawn"	CBR-II	5 (Pattern B)	Not Rec.	3 1/4	3 3/8	3 1/2
Cyclone	CB4000	4	3 1/2	3 3/4	3 7/8	4
"Spyker"	24-22	5	3	3.2	3.4	3.5
Andersons	"SS-2" Drop	NA	3 3/4	4	4 1/4	4 3/4

Note: These specified spreader settings should be used as starting points when calibrating a spreader to achieve the desired application rate of Acelepryn G. After calibration, regularly observe the amount of Acelepryn G that is applied to a known area and make further adjustments, as necessary, to maintain the appropriate application rate. Calibrate spreaders periodically to ensure that equipment wear or other factors have not altered the flow rate of Acelepryn G. Calibration instructions are included in the Owner's Manual that was provided by the equipment manufacturer at the time of purchase. Additional calibration information is available from your local Cooperative Extension Service.

CLEANING APPLICATION EQUIPMENT

Thoroughly clean all application equipment after use to reduce the risk of forming hardened deposits that might become difficult to remove. Thoroughly rinse application equipment with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean equipment near wells, or other water sources. Dispose of waste rinse water in accordance with local regulations.

INTEGRATED PEST MANAGEMENT (IPM) PROGRAMS

Acelepryn G is recommended for IPM programs on turf and landscape ornamentals because it does not directly impact natural arthropod predator and parasitoid populations including ladybird beetles, lacewings, minute pirate bugs and predatory mites. The feeding behavior of beneficial arthropods will aid in extending natural control of other insect and mite pests and will reduce the possibility of secondary pest outbreaks. Acelepryn G will reduce the target pest species that may serve as a food source for beneficial arthropods.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to products used repeatedly for control. When this occurs, the labeled dosages fail to suppress the pest population below threshold levels. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of active classes of insecticides on succeeding generations and targeting the most susceptible life stage. Consult your local Syngenta representative, Cooperative Extension Service specialist or pest control advisor for the latest information on resistance management in your area.

RESTRICTIONS

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production.
- Do not formulate this product into other end-use products without written permission from Syngenta.
- Do not apply this product in commercial nurseries and greenhouses.
- Do not apply more than 250 pounds (equivalent to 0.5 lb of active ingredient) of product per acre per year in broadcast applications to turfgrass.
- Do not apply more than 100 pounds (equivalent to 0.2 lb of active ingredient) of product per acre in a single application on sod farms.
- Wait a minimum of 7 days to retreat

Acelepryn® G

- Do not apply by air.
- The following restrictions are required to permit use of Acelepryn G in the State of New York:

Golf courses:

Greens, tee boxes: Do not apply this product within 25 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

Fairways: Do not apply this product within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

All other use sites:

Do not apply this product within 100 feet of a water body (lake, pond, river, stream, wetland or drainage ditch).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Not for storage in or around the home.

Pesticide Disposal

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product must be disposed of on site or an approved waste disposal facility.

Container Handling

Nonrefillable container. Do not reuse or refill this container. Completely empty plastic bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then offer for recycling, if available or dispose of empty plastic bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Acelepryn® G

Acelepryn®, the ALLIANCE FRAME 
the SYNGENTA Logo 
and the PURPOSE ICON 
are Trademarks of a Syngenta Group Company

©2014 Syngenta

"Vicon" is a registered trademark of Kverneland Group

"Lely" is a registered trademark of Maasland N.V.

"Scotts" is a registered trademark of OMS Investments, Inc.

"LESCO" is a registered trademark of LESCO Technologies, L.L.C.

"Prizelawn" is a registered trademark of White Castle System, Inc.

"Spyker" is a registered trademark of Spyker Spreaders, L.L.C.

"Accupro 2000" and "SS-2" Drop are trademarks of The Andersons, Inc.

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1500A-L1B 0414
4040317